

Infinite Characteristic Length in Small-World Systems

Cristian F. Moukarzel and Marcio Argollo de Menezes

Instituto de Física, Universidade Federal Fluminense, CEP 24210-340, Niterói, RJ, Brazil

It was recently claimed that on d -dimensional small-world networks with a density p of shortcuts, the typical separation $s(p) \sim p^{-1/d}$ between shortcut-ends is a characteristic length for shortest-paths (M. E. J. Newman and D. J. Watts, “Scaling and percolation in the small-world network model” *cond-mat/9904419*). This contradicts an earlier argument suggesting that no finite characteristic length can be defined for bilocal observables on these systems (M. Argollo de Menezes, C. Moukarzel and T. J. P. Penna, “First-order transition in small-world networks”, *cond-mat/9903426*). We show analytically, and confirm by numerical simulation, that shortest-path lengths $\ell(r)$ behave as $\ell(r) \sim r$ for $r < r_c$ and as $\ell(r) \sim r_c$ for $r > r_c$, where r is the Euclidean separation between two points and $r_c(p, L) \sim p^{-1/d} \log(L^d p)$. This shows that the mean separation s between shortcut-ends is *not* a relevant length-scale for shortest-paths. The true characteristic length $r_c(p, L)$ *diverges* with system size L no matter the value of p . Therefore no finite characteristic length can be defined for small-world networks in the thermodynamic limit.

PACS numbers: 05.10.-a, 05.40.-a, 05.50.+q, 87.18.Sn

The recent observation [1] that a non-extensive amount of long-range bonds, or “shortcuts”, on a regular d -dimensional lattice suffices to drastically shorten shortest-path distances, has triggered appreciable interest in these so-called “small-world” networks [2–9].

Small-world networks consist of a regular d -dimensional lattice of linear size L to which $L^d p$ long-ranged bonds have been added, connecting randomly chosen pairs of sites. Several recent studies have concentrated on the behavior of the average shortest-path distance $\bar{\ell} = 1/N^2 \sum_{\langle ij \rangle} \ell_{ij} <$, where ℓ_{ij} is the minimum number of links that must be traversed to join sites i and j , and $\langle \rangle$ means average over disorder realizations. On a regular lattice one has $\bar{\ell} \sim L$, while on a random graph of L^d sites, $\bar{\ell} \sim \log(L)$. On small-world networks, which interpolate between these two limits, one finds [1,3,6] $\bar{\ell} \sim L$ if $L^d p \ll 1$, but $\bar{\ell} \sim \log L$ if $L^d p \gg 1$. Thus on a large system a small density of shortcuts is enough to have shortest-path distances which are characteristic of random graphs.

It has been suggested [3] that for any fixed density p of long-ranged bonds, a *crossover-size* $L^*(p)$ exists, above which the average shortest-path distance $\bar{\ell}$ increases only logarithmically with L . Renormalization Group arguments [6] as well as numerical measurements [4,6,8] indicate that this crossover length diverges as $p \rightarrow 0$ as $L^*(p) \sim p^{-1/d}$. While some authors see this result as evidence of a continuous phase transition at $p = 0$ [6,9], it

has also been argued [8] that no finite correlation length can be defined on these systems, and thus that the small-world transition is first-order.

Clearly, the normalized shortest-path distance $\mathcal{L}(p) = \bar{\ell}/L$ undergoes a discontinuity at $p = 0$ in the thermodynamic limit. Thus, the “small-world” transition is technically a *first-order*, or discontinuous, transition [8]. A subtler and still controversial point [6,8,9] is whether it is possible to identify a correlation length $\xi(p)$ diverging at $p = 0$, i.e. if the small-world transition is associated with some sort of *critical behavior*. This would mean that $p = 0$ is a *first-order critical point* (FOCP) [11]. As discussed by Fisher and Berker [11], a FOCP is a critical point where the thermal eigen-exponent attains its maximum possible value $y_t = d$, thus allowing the *coexistence* of two different critical phases. Consequently at a FOCP one has $\nu = 1/y_t = 1/d$, and some finite-size corrections will be dictated by this exponent.

A FOCP appears for example in one-dimensional percolation [10], where the order parameter P_∞ (the density of the spanning cluster) is discontinuous at $p_c = 1$, while the correlation length ξ diverges as $p \rightarrow 1$ as $\xi \sim (1 - p)^{-1}$. Finite-size effects as e.g. the width of the critical region are of order L^{-1} . Because of this, P_∞ is non-zero even for $p \neq 1$, if $L < \xi \sim (1 - p)^{-1}$. Another example of a FOCP is the Ising model in one dimension, where the magnetization density $\langle m \rangle (T)$ is discontinuous at $T = 0$ while the correlation length ξ diverges as $\xi \sim e^{J/T}$.

At a normal first-order point, on the other hand, no critical behavior exists (i.e. all relevant L -independent length-scales remain finite), but the eigen-exponent d also shows up in finite-size corrections. In order to illustrate this, let us consider [11] a field-driven first-order transition at $h = 0$ (like e.g. in the two-dimensional Ising model below T_c). Assume that the central spin in a system of linear size L is pinned in the direction opposite to the external field h , and let us take for simplicity $T \sim 0$. If L is large enough, the bulk energy associated with h will be predominant, and $m(L, h)$ will have the same sign as h . On the other hand if L is small, it will be energetically favorable not to break any bonds and thus all spins will point in the $-h$ direction. The crossover length L^* at which magnetization inversion happens can be estimated by equating the total field contribution $L^d h$ to the energy cost associated with the breaking a finite number of bonds around the pinned spin, and one obtains $L^* \sim h^{-1/d}$ [11]. So we see that there may be a diverging crossover size in a first-order transition, although there is no diverging correlation length in this case.

different shortcuts. Notice that $V(t) \sim t$ for $t \ll s$ (only the primary sphere is important) while later $V(t) \sim e^{t/s}$ when $t > s$ (proliferation of secondary spheres). Because of this change in the behavior of $V(t)$ at $t \approx s$, Newman and Watts [9] suggest that s is the equivalent of a correlation length.

The total infected volume $V(t)$ in (1) is the sum of the primary volume $V_1(t)$ plus the secondary volume $V_2(t)$. The primary volume is simply $V_1(t) = t^d \Omega_d$, where Ω_d is the volume of a d -dimensional hypersphere of unit radius. According to our discussion above, we are interested in calculating the probability $P_S(r)$ for an individual located at a distance r from A to become infected by a secondary sphere before being hit by the primary sphere. Because the secondary volume $V_2(t)$ is *randomly distributed* in the total available space L^d , it is easy to calculate the probability $p_2(t)$ for an individual to become infected by the secondary version of the disease at time t or earlier. This is simply the fraction of the total volume that is covered by the secondary volume: $p_2(t) = V_2(t)/L^d$, and does not depend on the Euclidean distance from the primary source at A . If the individual does not become immunized by the secondary disease first, it will certainly be killed by the primary infection at time $t = d_E(A, x)$, when the primary sphere reaches him. Thus the relevant quantity is $p_2(t)$ evaluated at $t = d_E(A, x)$, which measures the probability $P_S(d_E(A, x))$ for an individual at x to survive. In one dimension one obtains, using (1) and $V_1(t) = 2t$,

$$P_S(d_E(A, x)) = \frac{s}{2L} \left(e^{4d_E(A, x)/s} - \frac{4d_E(A, x)}{s} - 1 \right), \quad (2)$$

which goes to zero when $L \rightarrow \infty$ no matter how large $d_E(A, x)$ is. Thus there is no *finite* safety distance in the thermodynamic limit, meaning that r_c diverges with system size. The size-dependence of the “safety distance” r_c can be estimated from (2) by equating $P_S \approx \text{constant}$, from which one obtains $r_c \sim s \ln(L/s)$.

These heuristic arguments are confirmed by an exact calculation of $\ell(r)$ [13], as described in the following. Newman and Watts equation for $V(t)$ [9] can be exactly solved in all dimensions, and using this solution it is possible to calculate the average shortest-path distance $\ell(r)$ between two points separated by an Euclidean distance r . The result is [13]:

$$\ell(r) \sim \begin{cases} r & \text{for } r < r_c = p^{-1/d} \log(p^{1/d} L) \\ r_c & \text{for } r > r_c \end{cases} \quad (3)$$

We confirmed this result numerically in one dimension, by measuring the average shortest path distance $\ell(r)$ as a function of Euclidean separation r , for several values of L and p . Equation (3) then means that the rescaled shortest-path distance $\tilde{\ell}(r) = \ell(r)/r_c$ should be a universal function of the rescaled Euclidean distance $\tilde{r} = r/r_c$.

This is confirmed by the fact that all data displayed in Fig. 2 fall on a single curve.

Notice that the characteristic length $r_c \sim s \log(L/s)$ is much larger than s , the mean separation between shortcuts. The reason why s is not a relevant length scale for shortest-path distances is not difficult to understand. Consider the shortest-path distance $\ell(A, x)$ between A and some point x as in Fig. 1. While the average distance one has to travel from A in order to find a shortcut end is s , this shortcut has a typical length $\sim L$ and thus will take us too far away from A to be useful in shortening the distance to x .




FIG. 2. Rescaled shortest path distance $\ell(r)/r_c$ versus rescaled Euclidean distance r/r_c , for one-dimensional small-world networks. The good collapse of these data onto a single curve clearly confirm Eq. (3) in the text. The characteristic length r_c for shortest-paths behaves as $p^{-1/d} \log(L^d p)$ and thus diverges, for all p , with system size L .

According to (3), only when $d_E(A, x)$ is larger than $r_c = s \log(L^d p)$ is it more “convenient” to go from A to x through shortcuts than directly through the lattice. This result means that the typical shortest path in the regime of long distances $r \gg r_c$ contains $\mathcal{O}(\log(L^d p))$ shortcuts, since the typical “price” paid for each shortcut traversed is the average distance between shortcut ends, i.e. s .

Thus s is *not* a relevant scale for shortest-path lengths $\ell(r)$ and therefore it does not play the role of a correlation length as suggested recently [9]. A physically more appealing interpretation of s is to say [9] that it is a diverging *timescale* for the dynamic process of disease spreading.

As might be clear by now, the impossibility to define a finite correlation length is a consequence of the fact that the locations of the secondary spheres are uncorrelated with the location of the primary infection. In other

words, while the typical separation between ends of different shortcuts $s(p)$ is finite for $L \rightarrow \infty$, the typical separation between both ends of the same shortcut, which is an important quantity for shortest-paths, scales as L . As a consequence of this, the number of shortcuts with both ends inside a region of any finite size ξ goes to zero when $L \rightarrow \infty$, which already implies the impossibility to define a finite correlation length [8]. Thus, although our demonstration here that no finite correlation length can be defined for shortest-path lengths is more rigorous, the same conclusion can be reached by using simple arguments [8].

A different situation would certainly arise if shortcuts had a length-dependent distribution, for example if each site i is connected with probability p to another site j chosen with probability $r^{-\alpha}$, where r is the distance between i and j and α is a free parameter. For $\alpha \rightarrow 0$, this model is the same as discussed here, while for α large enough one would only have short-range connections and thus there would be no logarithmic regime in $\bar{\ell}$, even for $p = 1$. We speculate that the transition in p could be shifted to a finite p_c for some intermediate α values. We are presently working on this extended model [12].

ACKNOWLEDGMENTS

We acknowledge useful discussions with P. M. C. de Oliveira, T. J. P. Penna, D. J. Watts, and M. Newman. This work is supported by CAPES and FAPERJ.

- [10] *Introduction to Percolation Theory*, D. Stauffer and A. Aharony, 2nd ed. (Taylor & Francis 1994).
- [11] M. E. Fisher and A. N. Berker, Phys. Rev. B **26** (1982), 2507.
- [12] M. Argollo de Menezes, C. Moukarzel and T. J. P. Penna, in preparation.
- [13] C. Moukarzel, 1999, to be published.

-
- [1] D. J. Watts and S. H. Strogatz, "Collective dynamics of small-world networks", Nature **393**, 440 (1998).
 - [2] H. Herzel, "How to quantify "small world" networks?", Fractals **6**, 4 (1998).
 - [3] M. Barthélemy and L.A.N. Amaral, "Small-world networks: evidence for a crossover picture", *cond-mat/9903108*, Phys. Rev. Lett. to appear.
 - [4] A. Barrat, "Comment on 'Small-world networks: evidence for a crossover picture' ", *cond-mat/9903323*.
 - [5] R. Monasson, "Diffusion, localization and dispersion relations on small-world lattices", *cond-mat/9903347*.
 - [6] M. E. J. Newman and D. J. Watts, "Renormalization group analysis of the small-world network model", *cond-mat/9903357*.
 - [7] A. Barrat and M. Weigt, "On the properties of small-world network models", *cond-mat/9903357*.
 - [8] M. Argollo de Menezes, C. Moukarzel and T. J. P. Penna, "First-order transition in small-world networks", *cond-mat/9903426*.
 - [9] M. E. J. Newman and D. J. Watts, "Scaling and percolation in the small-world network model", *cond-mat/9904419*.